



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/82, 15/11, 15/55 C12N 15/56, A01H 5/00, 5/08 A01H 5/10	A1	(11) International Publication Number: WO 93/23551 (43) International Publication Date: 25 November 1993 (25.11.93)
(21) International Application Number: PCT/GB93/00979 (22) International Filing Date: 13 May 1993 (13.05.93) (30) Priority data: 9210273.0 13 May 1992 (13.05.92) GB (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; Imperial Chemical House, 9 Millbank, London SW1P 3JF (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : BIRD, Colin, Roger [GB/GB]; 31 Fairfax, Bracknell, Berkshire RG12 1YT (GB). FRAY, Rupert, George [GB/GB]; 60 Derby Grove, Lenton, Nottingham NG7 1PF (GB). GRIERSON, Donald [GB/GB]; 55 Ashby Road, Shepshed, Loughborough LE12 9BS (GB). SCHUCH, Wolfgang, Walter [DE/GB]; 14 Greenfinch Close, Heathlake Park, Crowthorne, Berkshire RG11 6TZ (GB). SEYMOUR, Graham, Barron [GB/GB]; 38 The Plantation, Worthing, West Sussex BN13 2AJ (GB). TUCKER, Gregory, Alan [GB/GB]; 34 Lansdowne Road, Shepshed, Loughborough LE12 9RS (GB).		(74) Agent: ROBERTS, Timothy, Wace; Zeneca Limited, c/o ICI Group Patent Services Department, P.O. Box 6, Bessmer Road, Welwyn Garden City, Herts AL7 1HD (GB). (81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: CO-ORDINATED INHIBITION OF PLANT GENE EXPRESSION (57) Abstract <p>Process for the inhibition of two or more target genes which comprises introducing into the plant a single control gene which has distinct DNA regions homologous to each of the target genes and a promoter operative in plants adapted to transcribe from such distinct regions RNA that inhibits expression of each of the target genes. Constructs suitable for use in the process, as well as cells and plants containing such constructs are disclosed. Specific examples relate to the pectinesterase and polygalacturonase genes.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

- 1 -

CO-ORDINATED INHIBITION OF PLANT GENE EXPRESSION

This invention relates to novel DNA constructs, plant cells containing the constructs and plants derived therefrom. In particular it involves the use of recombinant DNA technology to control, and more specifically to inhibit, the expression of two or more genes in plants.

Plant development is a complex physiological and biochemical process requiring the co-ordinated expression of many genes. The production of new plant varieties with improved agricultural or commercial qualities can be achieved by modifying this co-ordinated pattern of gene expression. Such modifications have been achieved by conventional plant breeding techniques. However, the exact changes in gene expression that result in the production of the improved variety have not been readily characterised. More recently, recombinant DNA techniques have been used to modify the expression patterns of individual, specific plant genes without directly affecting the expression of other plant genes. In this way, the expression pattern of an individual gene can be either enhanced or inhibited either in the whole plant or in specific tissues or developmental stages.

The inhibition of specific individual plant genes has been achieved by the introduction into the plant of novel genes designed to express RNA homologous, in part, to the endogenous plant gene. In several cases, it has been demonstrated that expression of the target gene can be inhibited by two different strategies. These involve the introduction of specific genes designed to express either antisense or sense RNA. A typical example is the down-regulation of the gene encoding the tomato fruit cell wall

- 2 -

enzyme, polygalacturonase, by the expression of either antisense RNA (Smith et al 1988 Nature 344, 724-726) or sense RNA (Smith et al 1990 Mol Gen Genet 224, 477-481). A further example is the down-regulation of the gene encoding chalcone synthase in petunia by either sense or antisense RNA.

The mechanisms by which the expression of a specific gene is inhibited by either antisense or sense RNA genes are not clearly understood. It has been proposed that RNA-RNA duplex molecules may be formed within the cells resulting in the inhibition of expression. However, other and perhaps different mechanisms may operate for the two strategies for down-regulation. Specific individual genes have been inhibited by greater than 99% by the two strategies independently.

According to the present invention, we provide a process for the inhibition of two or more target genes which comprises introducing into the plant a single control gene which has distinct DNA regions homologous to each of the target genes and a promoter operative in plants adapted to transcribe from such distinct regions RNA that inhibits expression of the target genes. The distinct DNA regions homologous to each of the target genes may be either sense or antisense strands. This invention can be used to generate plants with the combined benefits of down-regulation of several individual genes or families of related genes.

The present invention further provides DNA constructs containing a plant promoter positioned to transcribe an RNA strand from at least two distinct DNA regions homologous to DNA from each of at least two target genes. The distinct

- 3 -

DNA regions may be joined sequentially or separated by a spacer region (preferably relatively short) provided such spacer region does not contain a transcription stop signal. The RNA transcribed from this single gene will contain regions homologous to the RNA transcribed from the two or more target genes.

The present invention further comprises novel cells and plants adapted to carry out the process of the invention, or which (or ancestors of which) have been transformed with the constructs of the invention.

The use of DNA constructs according to the invention offers several advantages over alternative, more complex methods of generating plants expressing homologous RNA to two or more individual target genes:

1. Plants can be generated from a single round of transformation thereby eliminating the extra time and complexity of multiple rounds of transformation.
2. The use of a single gene will ensure that RNA homologous to the two or more target genes is transcribed at the same rate. Otherwise, multiple individual homologous genes located in a single construct may have different transcription rates in transformed plants.
3. The segregation patterns of a single, multiple-function gene in progeny of a transformed plant will be simpler than the segregation patterns of multiple, single-function genes that have been combined in the same plant, either by sexual crossing or multiple transformation. This will have significant benefits for subsequent plant breeding.

- 4 -

The genes to which the present invention can be applied include all plant genes for which there is an advantage in down-regulating their activity. In particular the invention can be applied to genes involved in fruit development or ripening-related processes of commercially important fruit-bearing plants, in particular tomato. These could involve combinations of any of the genes encoding: cell wall hydrolases (eg. polygalacturonase, pectin esterase, β -(1-4)glucanase (cellulase), β -galactanase (β -galactosidase)); ethylene biosynthetic enzymes (eg 1-aminocyclopropane-1-carboxylate synthase, ethylene-forming enzyme); carotenoid biosynthetic enzymes (eg prephytoene or phytoene synthase, phytoene desaturase). In addition to genes encoding known enzymes, other genes (eg those showing homology to: pTOM36, pTOM38, pTOM66, pTOM75, pTOM99, pTOM136) with ripening enhanced expression have been identified and may be used in combination with any of the other genes.

DNA constructs according to the invention preferably comprise a base sequence at least 10 bases in length for each of the unrelated target genes for transcription into RNA. There is no theoretical upper limit to the base sequence for each homologous sequence - it may be as long as the relevant mRNA produced by the cell - but for convenience it will generally be found suitable to use sequences between 100 and 1000 bases in length. There is no theoretical limit to the length of the region separating the homologous sequences. The preparation of such constructs is described in more detail below.

The preferred DNA for use in the present invention is DNA derived from cDNA or genomic DNA of the target genes.

- 5 -

The required homologous DNA can be obtained in several ways: by cutting with restriction enzymes an appropriate sequence of such DNA; by synthesising a DNA fragment using synthetic oligonucleotides which are annealed and then ligated together in such a way as to give suitable restriction sites at each end; by using synthetic oligonucleotides in a polymerase chain reaction (PCR) to generate the required fragment with suitable restriction sites at each end. The DNA fragments from each of the unrelated target genes is then either ligated together prior to cloning or cloned sequentially into a vector containing upstream promoter and downstream terminator sequences.

Recombinant DNA and vectors according to the present invention may be made as follows. Suitable vectors containing the desired base sequences for transcription are treated with restriction enzymes to cut the sequences out. The DNA strands so obtained are cloned either simultaneously or sequentially into a second vector containing the desired promoter sequence (for example cauliflower mosaic virus 35S RNA promoter or the tomato polygalacturonase gene promoter sequence - Bird et al., Plant Molecular Biology, 11, 651- 662, 1988) and the desired terminator sequence (for example the 3' of the Agrobacterium tumefaciens nopaline synthase gene, the nos 3' end).

According to the invention we propose to use both constitutive promoters (such as cauliflower mosaic virus 35S RNA) and inducible or developmentally regulated promoters (such as the ripe-fruit-specific polygalacturonase promoter) as circumstances require. Use of a constitutive promoter will tend to affect functions in all parts of the plant: while by using a tissue specific

promoter, functions may be controlled more selectively. Thus in applying the invention, e.g. to tomatoes, it may be found convenient to use the promoter of the PG gene (Bird et al, 1988, cited above). Use of this promoter, at least in tomatoes, has the advantage that the production of homologous RNA is under the control of a ripening-specific promoter. Thus the homologous RNA is only produced in the organ in which its action is required. Other ripening-specific promoters that could be used include the E8 promoter (Diekman & Fischer, 1988 cited above).

Vectors according to the invention may be used to transform plants as desired, to make plants according to the invention. Dicotyledonous plants, such as tomato and melon, may be transformed by Agrobacterium Ti plasmid technology, for example as described by Bevan (1984) Nucleic Acid Research, 12, 8711-8721. The process may also be adapted for use with other techniques for generating transgenic plants. Such transformed plants may be reproduced sexually, or by cell or tissue culture.

The degree of production of homologous RNA in the plant cells can be controlled by suitable choice of promoter sequences, or by selecting the number of copies, or the site of integration, of the DNA sequences according to the invention that are introduced into the plant genome. In this way it may be possible to modify expression of the target genes to a greater or lesser extent.

The constructs of our invention may be used to transform cells of both monocotyledonous and dicotyledonous plants in various ways known to the art. In many cases such plant cells (particularly when they are cells of dicotyledonous plants) may be cultured to regenerate whole

- 7 -

plants which subsequently reproduce to give successive generations of genetically modified plants. Examples of genetically modified plants according to the present invention include, as well as tomatoes, fruiting plants such as mangoes, peaches, apples, pears, strawberries, bananas and melons.

The specific embodiment of the invention that we have so far studied most thoroughly is the inhibition of both polygalacturonase and pectin esterase in ripening tomatoes. Modified tomato plants have been produced which contain a novel gene designed to express RNA with regions individually homologous to the tomato PG and PE genes. Fruit from some of the primary transformants have significant reductions of both PG and PE activity in the ripening fruit. This reduced enzyme activity is stably inherited as a single gene in progeny of the primary transformants. Plants such as these exhibit the combined benefits of reduced PG and PE activity and will be useful in the production of tomatoes with improved quality for fresh market and processed fruit.

The invention will now be described further with reference to the accompanying drawings, in which:

Figure 1 shows the structure of the PG-PE sense gene in pPGPE;

Figure 2 is a diagram of the strategy for construction of the PG-PE antisense gene in pJF1;

Figure 3 is a graph of PG and PE activities in fruit from a tomato plant transformed with pPGPE ("dpp" = "days post breaker");

Figure 4 shows PG and PE activity in fruit from 28 tomato plants transformed with a PE-PG sense construct;

Figure 5 shows PG and PE activity in fruit from 32 tomato plants transformed with a PE-PG antisense construct.

- 8 -

Example 1

Construction of combined PG-PE sense gene

A PG-PE sense gene was constructed by cloning bases 19 to 263 of the PG cDNA (pTOM6 - Grierson et al 1986 Nuc. Acids Res. 14, 8595- 8603) and bases 345 to 1665 of the PE cDNA (pPE1 - Ray et al 1988 Eur. J. Biochem 174, 119-124) into the multiple cloning site of the vector pDH51. pDH51 is a pUC based vector which contains a multiple cloning site between the CaMV 35S promoter sequence and the CaMV 3' polyadenylation sequence. The orientation of the PG and PE sequences was determined by restriction mapping and DNA sequence determination of the borders of the inserted fragments. After verification of the structure of the vector, the expression module was isolated by digestion with pvuII and transferred to the plant transformation vector Bin19 (Bevan 1984 Nuc. Acids Res. 12, 8711-8721) to give the vector pPGPE (also known as pSB1).

Example 2

Construction of combined PG-PE antisense gene

A PG-PE antisense vector (PJF1) was constructed as shown in figure 2. A 686 bp EcoRI-XbaI fragment from pPBI that contained the CaMV 35S promoter sequence and the first 150 bases from the 5' end of the PG cDNA (pTOM6) was isolated. The fragment was cloned into the EcoRI-XbaI sites of the PE antisense vector pJR101ARi after removal of the CaMV 35S promoter sequence. The construction of the vector was confirmed by PCR and DNA sequence analysis.

Example 3

- 9 -

Generation of transformed tomato plants

Vectors from Example 1 were transferred to Agrobacterium tumefaciens LBA4404 (a micro-organism widely available to plant biotechnologists) and were used to transform tomato plants. Transformation of tomato stem segments followed standard protocols (e.g. Bird et al Plant Molecular Biology 11, 651-662, 1988). Transformed plants were identified by their ability to grow on media containing the antibiotic kanamycin. Plants were regenerated, grown to maturity and the presence of the PG-PE sense gene was confirmed by genomic Southern analysis.

Example 4

Analysis of fruit from transformed plants

The PG and PE activities were assayed in ripening fruit from 3 plants transformed with pPGPE and several un-modified plants. The PG and PE activity was reduced by various extents in the fruit from all three transformed plants. The results for fruit from one of the transformed plants are shown in Figure 3.

Example 5

Analysis of progeny of transformed plants

Self-fertilised seed from the pPGPE-transformed plant with the greatest reductions in PG and PE activities were germinated in the presence of kanamycin. Plants that were resistant to kanamycin were grown to maturity and enzyme activities in the fruit were analysed. These plants also

- 10 -

had reduced PG and PE activity. Thus, the inheritance of the reduced PG and PE phenotype was confirmed.

EXAMPLE 6

The sense construct pSB1 prepared in Example 2 was used to transform tomatoes of the variety UC82B, using the same protocol as in Example 3. Transformants were grown in the glasshouse. PG and PE enzyme activities were assayed in ripe fruit. Fruit from the first 28 plants to mature were analysed. The results are presented graphically in Figure 4, ranked in order of decreasing PG level. These results confirm that plants with substantial reductions in both PG and PE activity were obtained from the population of transformants. In all plants with significantly reduced activity of one enzyme, the other was also reduced. Thus the sense down-regulation of the two genes was tightly linked.

EXAMPLE 7

Expression of genes in plants containing sense constructs.

Selfed progeny were grown from two primary transformants obtained from Example 3 with single sites of insertion (E148C11 - PG expression 10% that of normal fruit, PE expression 20% normal; E143C44 - PG expression 5% normal, PE expression 16% normal). Stable inheritance of the double sense gene was confirmed by genomic Southern analysis. Homozygous and azygous progeny were identified. Assays of ripe fruit PG and PE activities confirmed that the low PG and PE phenotype segregated with the sense gene:

PG activity	PE activity
-------------	-------------

- 11 -

E143C44.16	Homozygous	1	108
E143C44.30	Azygous	34	307
E148C11.1	Homozygous	2.5	98
E148C11.25	Azygous	34	480

(all nmol/hr/ μ g protein)

EXAMPLE 8

Expression of genes in plants containing antisense constructs

68 Transformants with pJF1 (from Example 2) in were generated in the tomato variety UC82B. PG and PE enzyme activities were assayed in ripe fruit. Fruit from the first 32 plants to mature were analysed. Untransformed UC82B is included as a control. The results are presented graphically in Figure 5, ranked in order of decreasing PG level. As with the sense plants, reduced activities of the two enzymes were tightly linked.

This data confirms the utility of the antisense construct pJF1 (Example 2).

Analysis of selfed progeny confirms that both the antisense gene and the reduced PG/PE phenotype are inherited.

- 12 -

We claim:

1. A process for the inhibition of two or more target genes which comprises introducing into the plant a single control gene which has distinct DNA regions homologous to each of the target genes and a promoter operative in plants adapted to transcribe from such distinct regions RNA that inhibits expression of each of the target genes.
2. A process as claimed in claim 1 in which at least one of the DNA regions homologous to a target gene is adapted to produce sense RNA.
3. A process as claimed in claim 1 in which at least one of the DNA regions homologous to a target gene is adapted to produce antisense RNA.
4. A process as claimed in claim 1 in which the promoter is a constitutive promoter.
5. A process as claimed in claim 1 in which the promoter is an inducible promoter.
6. DNA constructs useful in the process of claim 1 containing a plant promoter positioned to transcribe an RNA strand from at least two distinct DNA regions homologous to DNA from each of at least two target genes.
7. A DNA construct as claimed in claim 6 in which at least one of the DNA regions homologous to a target gene is adapted to produce sense RNA.
8. A DNA construct as claimed in claim 6 in which at least one of the DNA regions homologous to a target gene is

- 13 -

adapted to produce antisense RNA.

9. A DNA construct as claimed in claim 6 in which the target genes are fruit ripening genes.

10. A DNA construct as claimed in claim 6 in which one of the target genes is a polygalacturonase or pectinesterase gene.

11. A plant cell transformed with a construct claimed in claim 6.

12. Plant cells claimed in claim 11 which are cells of tomato.

13. Plant cells claimed in claim 11 which are cells of mangoes, peaches, apples, pears, bananas, melons or strawberries.

14. Plants containing cells claimed in either of claims 11 and 12.

15. Fruit or seeds of plants claimed in either of claims 13 or 14.

16. Tomato seeds as claimed in claim 15.

17. Tomato fruit as claimed in claim 17.

FIG. 1

The Structure of the PG-PE sense gene in pPGPE




FIG. 2 (1/2)
Construction of PG-PE antisense gene in pJF1




FIG. 2 (2/2)

pJR101ARi






FIG. 3
PG and PE activities in fruit from a plant transformed with pPGPE




SUBSTITUTE SHEET

PG AND PE AVERAGE ACTIVITIES
FIG. 4 OF SENSE CONSTRUCT PLANTS



**FIG. 5 PG AND PE AVERAGE ACTIVITIES
OF ANTISENSE CONSTRUCT PLANTS**



INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 93/00979

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/82; A01H5/00;	C12N15/11; A01H5/08;	C12N15/55; A01H5/10 C12N15/56
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12N ; A01H	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	EP,A,0 455 316 (INSTITUT GENE BIOLOGISCHE FORSCHUNG BERLIN) 6 November 1991	1,3,4,6, 8,11,14
Y	see page 15, line 29 - page 16, line 11	5,9,10, 12,15-17
Y	EP,A,0 341 885 (ICI) 15 November 1989	2,5,7,9, 10,12, 15-17
Y	WO,A,9 012 084 (DNA PLANT TECHNOLOGY CORPORATION) 18 October 1990	2,7
A	WO,A,9 108 299 (ICI) 13 June 1991	2,7
	see the whole document	
	-/--	
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
26 AUGUST 1993	20.09.93	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	MADDOX A.D.	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claims No.
A	EP,A,0 271 988 (ICI) 22 June 1988 see the whole document -----	3-5,8-17
A	EP,A,0 416 572 (HOECHST) 13 March 1991 see claim 10 -----	1,10
O,A	J. CELL. BIOCHEM. SUPPL., KEYSTONE SYMPOSIUM, HELD JAN. 10-17, 1991. vol. 15A, 1991, page 81 TIEMAN, D.L., ET AL. 'Inhibition of pectin methylesterase (PME) gene expression in transgenic tomato fruit by anti-sense RNA' see abstract A353 -----	10

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 9300979
SA 73948

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

26/08/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0455316	06-11-91	DE-A- 4013144	24-10-91
EP-A-0341885	15-11-89	AU-B- 628818	24-09-92
		AU-A- 3469789	16-11-89
		JP-A- 2031625	01-02-90
WO-A-9012084	18-10-90	US-A- 5034323	23-07-91
		AU-A- 5412390	05-11-90
		EP-A- 0465572	15-01-92
		JP-T- 4504800	27-08-92
		WO-A- 9011682	18-10-90
		US-A- 5231020	27-07-93
WO-A-9108299	13-06-91	AU-A- 6974591	26-06-91
		EP-A- 0502995	16-09-92
		JP-T- 5504056	01-07-93
EP-A-0271988	22-06-88	AU-B- 633202	21-01-93
		AU-A- 7435091	11-07-91
		AU-A- 8095687	12-05-88
		EP-A- 0532060	17-03-93
		JP-A- 63164892	08-07-88
		US-A- 5073676	17-12-91
EP-A-0416572	13-03-91	DE-A- 3929741	28-03-91
		AU-B- 634575	25-02-93
		AU-A- 6218690	14-03-91
		CA-A- 2024720	08-03-91
		JP-A- 3108484	08-05-91